

*Rickets and lamellar ichthyosis.* 71

*Perineal erythema as first sign of Kawasaki syndrome.* 75

*Multiple subcutaneous nodules due to alveolar rhabdomyosarcoma.* 78

*Skin lesions in neonates. A study from India.* 83

*Laser treatment of anterior cervical hypertrichosis.* 89

*Evaluation of safety and efficacy of a galenic preparation of 1% propranolol in 89 cases of hemangioma.* 93

*Frontal whorl. An early sign of peripheral neurofibromatosis?* 113

*Segmental neurofibromatosis associated with hypopigmentation "en confetti".* 114

*Self-healing localized congenital dystrophic epidermolysis bullosa.* 115



Fig. 1: at 3 mo. onset of oral propranolol; Fig. 2: at 14 mo. propranolol stopped; Fig. 3: at 30 mo. rebound.

*Neuroid congenital melanocytic nevus with hemangioma.* 116

*Nevus spilus resembling congenital melanocytic nevus.* 117

*Late onset hairy nevus with perifollicular hypermelanosis.* 118

*Incomplete development of the hallux nail.* 119

*Congenital neurofibroma with hypertrichosis.* 120

*Rebound of hemangioma after oral propranolol.* 121

*Vascular disorders in metameric hemangioma.* 122

*Urticaria-angioedema on hemangioma.* 124

*Keratolysis exfoliativa.* 125

*Hyperpigmentation of the oral cavity type Laugier-Hunziker.* 126

*Dermatitis artefacta.* 127

*Bullous mastocytoma.* 128